

## Field Evaluation Of Certain Newer Insecticides Against Chilli Thrips, *Scirtothrips Dorsalis* (Hood)

K Vanisree, S.Upendhar, P Rajasekhar, G Ramachandra Rao And V Srinivasa Rao



### ABSTRACT

The field experiment conducted during *Kharif* 2008-09 and 2009-10 on evaluation of certain new insecticides indicated that spinosad 0.015 per cent was found most effective in reducing the population of *S. dorsalis* as well as in increasing yields. It attains highest cost benefit ratio followed by diafenthiuron 0.045%, pymetrozine 0.02% and fipronil 0.01%. Indoxacarb 0.015 % and flubendiamide 0.012 %. Further flubendiamide recorded the lowest mean per cent reduction over untreated check indicating their poor efficacy against *S. dorsalis* on chillies.

#### Keywords:

Chilli Thrips, Management, Chemical Control, Novel Insecticides.

#### Introduction

India has emerged today as the foremost producer and exporter of chillies contributing to almost 1/4<sup>th</sup> of the world's production. In India, chilli is grown in an area of 8.06 lakh ha, with a production of 12.98 lakh tonnes (Agricultural Statistics at a glance, 2009). The important chilli growing states in India are Andhra Pradesh, Orissa, Maharashtra, Karnataka and also in a number of other states as a round the year crop. In Andhra Pradesh, chilli is cultivated in an area of 1.89 lakh hectares with a production of 2.08 lakh tonnes. Guntur district in Andhra Pradesh alone contributes to over 35 per cent in area under chilli crop in India.

The important pests are thrips, *Scirtothrips dorsalis* (Hood), white mite, *Polyphagotarsonemus latus* (Banks), aphids, *Aphis gossypii* Glover and *Myzus persicae* Sulzer as sucking complex and tobacco caterpillar, *Spodoptera litura* (Fabricius) and pod borer, *Helicoverpa armigera* (Hubner) as pod borers (Rao and Ahmed, 1985). Chilli thrips, *Scirtothrips dorsalis* (Hood) (Thysanoptera : Thripidae) is a serious pest of *Capsicum annum* L. in India, responsible for leaf curling (Ananthkrishnan, 1971). It multiplies appreciably at a faster rate during dry weather periods and the yield loss caused by the thrips is reported to range from 30-90 per cent (Borah, 1987 and Varadharajan, 1994).

Guntur district in Andhra Pradesh is traditionally a chilli growing district with an area of 63,573 ha with high input usage under monocropping conditions. Further, intensive cultivation of input responsive high yielding varieties and hybrids and sole reliance on insecticides are the common features of chilli cultivation in Guntur district. The excessive dependence on insecticides, their over use and abuse has accelerated insect control problems through development of insecticide resistance (Reddy *et al.*, 1992), pest resurgence, pesticide residues (Joia *et al.*, 2001), reduction in natural enemy population and environmental contamination. Moreover, several of the chilli consignments meant for export were rejected stating higher insecticide residues being the culprit, thus lots of foreign exchange lost by way of rejections. Further there were several reports from farmers experiencing difficulties in pest control. Many conventional insecticides are being used to manage these pests with which many folds of resistance was reported in pests like *S. litura* (Prasad *et al.*, 2008), *Spodoptera exigua* (Hubner) (Wang *et al.*, 2002), *H. armigera* (Kranthi *et al.*, 2002) *etc.* The occurrence of insecticide resistance strains can be reduced or delayed by reducing the selection

K Vanisree, S.Upendhar, P Rajasekhar,  
G Ramachandra Rao And  
V Srinivasa Rao

From  
Department of Entomology,  
Agricultural College, Bapatla,  
Andhra Pradesh .

The Article Is Published On December  
2013 Issue & Available At

[www.scienceparks.in](http://www.scienceparks.in)

DOI: [10.9780/23218045/1202013/49](https://doi.org/10.9780/23218045/1202013/49)



pressure and by adopting insecticide resistance management strategies and alternate insecticides with novel mode of action. In view of the above constraints in chilli cultivation, it is felt high time to estimate the current status of insecticide resistance so as to corrolagate with field control problems besides evaluating newer insecticides with novel mode of action both under laboratory and field conditions so as to have better option on hand that could mitigate the present control failures and residue problems plausing the farming community.

**Materials and Methods**

The experiment was laid in a simple randomized block design with twelve treatments including untreated check, replicated thrice. Each plot measured 18m<sup>2</sup> and the gross area of the management trial was 800m<sup>2</sup>. Different group of insecticides fipronil (Phenyl pyrazole), Pymetrozine (Pyridine Azomethines), Imidacloprid (Neonecotinoid), Diafenthiuron (Thiourea), Vertimec and Emamectin benzoate (Avermectins), Spinosad (Spinosyn), Indoxacarb (Oxadiazine), Flubendiamide (Phthalic acid diamide), Clothianidin (Neonecotinoid) and Chlorfenapyr (pyrrole compound) was evaluated against *S.dorsalis*. A measured quantity of insecticidal solution/ powder was mixed with a little quantity of water and stirred well, after which the remaining quantity of water was added to obtain the required concentration of spray fluid. Sprayings were given by using a hand compression knapsack high volume sprayer during morning hours. The plot in each treatment was sprayed with respective insecticides ensuring uniform coverage of insecticide. The sprayer and the accessories were thoroughly washed before changing the insecticides and also rinsed with the spray fluid of the chemical to be applied next. The first spraying was given at 45 days after transplanting, when the incidence of thrips population was sufficiently built up in the experimental plots. A total of five sprays were given during the course of season at fortnightly interval. Observations on the pest incidence were recorded one day before the spraying as pre-treatment count and first, third, seventh and fourteenth day after spraying as the post- treatment counts. The population of both nymphs and adults of thrips were counted during early morning hours on terminal six leaves from 5 randomly selected plants in each plot and the plants were tagged leaving the border rows. The matured green pods were harvested four times in all treatments at 15 days interval during crop period and the cumulative yield of all the pickings were recorded plot wise and expressed in kg/ net plot. From this the yield per hectare was calculated. The per cent reduction in thrips population over control was calculated by modified abbott's formula followed by Flemming and Retnakaran (1985).

Percentage population reduction =

$$1 - \left( \frac{\text{Post treatment population in treatment}}{\text{Pre treatment population in treatment}} \times \frac{\text{Pre treatment population in untreated control}}{\text{Post treatment population in untreated control}} \right) \times 100$$

These percentages were transformed to the corresponding arc sine values and the data was subjected to statistical analysis. The yield data in each treatment was recorded separately and subjected to statistical analysis to test the significance of mean yield in different treatments.

The percent increase in yield over control in various treatments was calculated by using the following formula.

$$\text{Percent increase of yield in treatment over control} = \frac{\text{Yield in treatment} - \text{Yield in control}}{\text{Yield in control}} \times 100$$

**Results and Discussion****Field management studies****Kharif 2008-2009**

The results with regard to the mean efficacy of the treatments after the five rounds of application are presented in Table 1. The data recorded one day after imposition of treatments indicated that spinosad 0.015 per cent was the most effective and significantly superior over all other treatments by recording the highest reduction in population of 82.43 per cent over untreated check. The next best treatment was diafenthiuron 0.045 per cent with 81.26 per cent reduction in population over untreated check. Next best was pymetrozine 0.02 per cent with 78.64 per cent reduction in population and was significantly superior to the rest of the other treatments. The other treatment has shown good degree of control with more than 77 per cent mean reduction over control was fipronil 0.01 per cent (77.48 per cent) being significantly superior to remaining treatments. The other treatments that followed in the descending order of efficacy were imidacloprid 0.005 per cent, chlorfenapyr 0.01 per cent, clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003 per cent with 67.52, 59.38, 57.60, 56.29 and 52.61 per cent reduction in thrips population over untreated check, respectively. Among the treatments, indoxacarb 0.015 per cent (47.52 per cent) and flubendiamide 0.012 per cent (43.25 per cent) showed poor efficacy with less than 48 per cent reduction in population. However, all the treatments were significantly superior over control in bringing down the population of *S. dorsalis* on chillies during *kharif* 2008-09 at one day after spraying.

The data recorded at three days after treatment showed that the most effective treatment was spinosad 0.015 per cent with 93.49 per cent reduction in population of *S. dorsalis* over control and was significantly superior over all the other treatments. The next best treatment was diafenthiuron 0.045 per cent with 92.00 per cent reduction in population over untreated check. The treatments pymetrozine 0.02 per cent (87.32 per cent) and fipronil 0.01 per cent (86.67 per cent) being on par with more than 86 per cent reduction in population over untreated check. The other treatments that followed in the descending order of efficacy were imidacloprid 0.005 per cent, chlorfenapyr 0.01 per cent, clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003 per cent with 75.51, 67.42, 62.55, 59.32 and 57.60 per cent reduction in thrips population over untreated check, respectively. Among the treatments, indoxacarb 0.015 per cent (52.37 per cent) and flubendiamide 0.012 per cent (46.88 per cent) were found to be least effective with less than 53 per cent reduction in population over untreated check. However, all the treatments were effective and significant in reducing the population of *S. dorsalis* over control at three days after spraying.

The data recorded at seven days after five sprayings showed that spinosad 0.015 per cent was the most effective and significantly superior over all the other treatments by recording the highest reduction in population of 87.38 per cent over untreated check. The next best treatment was diafenthiuron 0.045 per cent with 85.23 per cent reduction in population and was significantly superior to the rest of the treatments. Closely behind was pymetrozine 0.02 per cent with 79.32 per cent reduction in population over untreated check. The treatment, fipronil 0.01 per cent came next and was also significantly superior over the rest with 77.52 per cent reduction in population over untreated check. The other treatments that followed in the descending order of efficacy were chlorfenapyr 0.01 per cent, imidacloprid 0.005 per cent, clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003 per cent with 69.56, 66.38, 60.55, 58.29 and 56.41 per cent reduction in thrips population over untreated check, respectively. Among the treatments, indoxacarb 0.015 per cent (50.46 per cent) and flubendiamide 0.012 per cent (45.45 per cent) were found least effective with less than 51 per cent reduction in population over untreated check. However, all the treatments were effective and significantly superior over the untreated control in bringing down the population of *S. dorsalis* at seven days after spraying.

The post treatment data at fourteen days after spraying showed that diafenthiuron 0.045 per cent was the most effective and significantly superior over all the other treatments with 68.17 per cent reduction in population over untreated check, which was on par with spinosad 0.015% with 67.69 per cent reduction in population over untreated check. The next best treatment was pymetrozine 0.02 per cent being significantly superior to the rest of the treatments with 64.22 per cent reduction over untreated check. The treatment, fipronil 0.01 per cent came next and was also significantly superior over the rest with 62.40 per cent reduction in population over untreated check. The other treatments that followed in the descending order of efficacy were imidacloprid 0.005 per cent, chlorfenapyr 0.01 per cent,

clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003 per cent with 47.53, 46.53, 42.13, 40.24 and 37.45 per cent reduction in thrips population over untreated check, respectively. Among the treatments, indoxacarb 0.015 per cent (35.53 per cent) and flubendiamide 0.012 per cent (32.90 per cent) were found to be the least effective with less than 36 per cent reduction in population over control. However, all the treatments were effective and significantly superior over the untreated control in bringing down the population of *S. dorsalis* at fourteen days after spraying.

The data pertaining to the overall mean efficacy of the treatments after the imposition of five sprays revealed that the most effective and the best treatment was spinosad 0.015 per cent with 82.75 per cent reduction in population of *S. dorsalis* over untreated check and was significantly superior over all the other treatments. The next best treatment was diafenthiuron 0.045 per cent with 81.66 per cent reduction in population over untreated check. Next came, pymetrozine 0.02 per cent with 77.38 per cent reduction in population and was significantly superior to the rest of the treatments. The other treatment that has shown good degree of control with more than 76 per cent reduction over control was fipronil 0.01 per cent (76.02 per cent) being significantly superior to remaining treatments. The other treatments that followed in the descending order of efficacy were imidacloprid 0.005 per cent, chlorfenapyr 0.01 per cent, clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003% with 64.24, 60.72, 55.71, 53.53 and 51.02 per cent reduction in thrips population over untreated check, respectively. Among the treatments, indoxacarb 0.015 per cent (46.47 per cent) and flubendiamide 0.012 per cent (42.12 per cent) were found least effective with less than 47 per cent reduction in population over untreated check. However, all the treatments were significantly superior over the untreated control in reducing the population of *S. dorsalis* on chillies after the five rounds of sprays during *kharif*2008-2009.

#### ***Kharif*2009-2010**

The results with regard to the mean efficacy of the treatments after the five rounds of application are presented in Table 2. The data recorded one day after imposition of treatments indicated that spinosad 0.015 per cent was the most effective and significantly superior over all other treatments by recording the highest reduction in population of 81.22 per cent over untreated control, which was on par with diafenthiuron 0.045 per cent with 80.06 per cent reduction in population and was significantly superior to the rest of the treatments. The next best treatments were pymetrozine 0.02 per cent (77.44 per cent) and fipronil 0.01 per cent (76.27 per cent) being on par with more than 76 per cent reduction in population over untreated check. The treatment, imidacloprid 0.005 per cent came next and was also significantly superior over the rest with 66.32 per cent reduction in population over untreated check. The treatments chlorfenapyr 0.01 per cent (58.18 per cent), clothianidin 0.003 per cent (56.40 per cent) and vertimec 0.001 per cent (55.08 per cent) being on par with more than 55 per cent mean reduction in population over untreated check. The treatments emamectin benzoate 0.003 per cent (51.41 per cent) and indoxacarb 0.015 per cent (46.32 per cent) were also significantly superior to the rest of the treatments with more than 46 per cent mean reduction in population over control. The remaining treatment, flubendiamide 0.012 per cent (42.05 per cent) showed poor efficacy with less than 43 per cent reduction in population. However, all the treatments were significantly superior over control in bringing down the population of *S. dorsalis* on chillies during *kharif* 2009-2010 at one day after spraying.

The data recorded at three days after treatment showed that the most effective treatment was spinosad 0.015 per cent with 92.81 per cent reduction in population of *S. dorsalis* over control and was significantly superior over all the other treatments. The next best treatment was diafenthiuron 0.045 per cent with 91.32 per cent reduction in population and was significantly superior to the rest of the treatments. The next best treatments were pymetrozine 0.02 per cent (86.64 per cent) and fipronil 0.01 per cent (85.99 per cent) being on par with more than 85 per cent reduction in population over untreated check. The other treatments that followed in the descending order of efficacy were imidacloprid 0.005 per cent, chlorfenapyr 0.01 per cent, clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003 per cent with 74.83, 66.74, 61.87, 58.64 and 56.92 per cent reduction in thrips population over untreated check, respectively. Among the treatments, indoxacarb 0.015 per cent (51.69 per cent) and flubendiamide 0.012 per cent (46.20 per cent) were found to be the least effective with less

than 52 per cent mean reduction in population over control. However, all the treatments were effective and significant in reducing the population of *S. dorsalis* over control at three days after spraying.

The post treatment data at seven days after five sprayings showed that spinosad 0.015 per cent was the most effective and significantly superior over all the other treatments by recording the highest reduction in population of 86.67 per cent over untreated control. The next best treatment was diafenthiuron 0.045 per cent with 84.52 per cent reduction in population and was significantly superior to the rest of the treatments. Closely behind were pymetrozine 0.02 per cent (78.61 per cent) and fipronil 0.01 per cent (76.81 per cent) being on par with more than 76 per cent reduction in population over untreated check. The other treatments that followed in the descending order of efficacy were chlorfenapyr 0.01 per cent, imidacloprid 0.005 per cent, clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003 per cent with 68.85, 65.67, 59.84, 57.58 and 55.70 per cent reduction in thrips population over untreated check, respectively. Among the treatments, indoxacarb 0.015 per cent (49.75 per cent) and flubendiamide 0.012 per cent (44.74 per cent) were found to be the least effective with less than 50 per cent mean reduction in population over untreated check. However, all the treatments were effective and significantly superior over the untreated control in bringing down the population of *S. dorsalis* at seven days after spraying.

The post treatment data at fourteen days after spraying showed that diafenthiuron 0.045 per cent was the most effective and significantly superior over all the other treatments with 67.55 per cent reduction in the population over untreated control, which was on par with spinosad 0.015 per cent with 67.07 per cent reduction in population and was significantly superior to the rest of the treatments. The next best treatments were pymetrozine 0.02 per cent (63.60 per cent) and fipronil 0.01 per cent (61.78 per cent) being on par with more than 61 per cent reduction in population over untreated check. The treatments imidacloprid 0.005 per cent (46.91 per cent) and chlorfenapyr 0.01 per cent (45.91 per cent) being on par with more than 45 per cent reduction in population and significantly superior over untreated check. The treatments clothianidin 0.003 per cent (41.51 per cent) and vertimec 0.001 per cent (39.62 per cent) being on par with more than 39 per cent reduction in population over untreated check. The treatments emamectin benzoate 0.003 per cent (36.83 per cent) and indoxacarb 0.015 per cent (34.91 per cent) being on par with a mean reduction in population of more than 34 per cent when compared to control. The remaining treatment, flubendiamide 0.012 per cent (32.28 per cent) showed poor efficacy with less than 33 per cent reduction in population over untreated control. However, all the treatments were effective and significantly superior over the untreated control in bringing down the population of *S. dorsalis* at fourteen days after spraying.

The data pertaining to the overall mean efficacy of the treatments after the imposition of five sprays revealed that the most effective treatment was spinosad 0.015 per cent with 81.94 per cent reduction in population of *S. dorsalis* over untreated control and was significantly superior over all the other treatments, which was on par with diafenthiuron 0.045 per cent with 80.86 per cent reduction in population and was significantly superior to the rest of the treatments. The next best treatments were pymetrozine 0.02 per cent (76.57 per cent) and fipronil 0.01 per cent (75.21 per cent) being on par with more than 75 per cent reduction in population over untreated check. The other treatments that followed in the descending order of efficacy were imidacloprid 0.005 per cent, chlorfenapyr 0.01 per cent, clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003 per cent with 63.43, 59.92, 54.90, 52.73 and 50.22 per cent reduction in thrips population over untreated check, respectively. Among the treatments, indoxacarb 0.015 per cent (45.67 per cent) and flubendiamide 0.012 per cent (41.32 per cent) were found to be the least effective with less than 46 per cent reduction in population over control. However, all the treatments were significantly superior over the untreated control in reducing the population of *S. dorsalis* on chillies after the five rounds of sprays during *khariif* 2009-2010.

#### **Overall cumulative mean efficacy of treatments during both the seasons**

The results with regard to the overall cumulative mean efficacy of the treatments during both the years are presented in Table 3. The mean data recorded one day after imposition of treatments indicated that spinosad 0.015 per cent was the most effective and significantly superior over all other treatments by recording the highest reduction in population of 81.83 per cent over untreated check, which was on par with diafenthiuron 0.045

per cent with 80.66 per cent reduction in population and was significantly superior to the rest of the treatments. The next best treatments were pymetrozine 0.02 per cent (78.04 per cent) and fipronil 0.01 per cent (76.87 per cent) being on par with more than 76 per cent reduction in population over untreated check. The other treatments that followed in the descending order of efficacy were imidacloprid 0.005 per cent, chlorfenapyr 0.01 per cent, clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003 per cent with 66.92, 58.78, 57.00, 55.69 and 52.01 per cent reduction in thrips population over untreated check, respectively. Among the treatments, indoxacarb 0.015 per cent (46.92 per cent) and flubendiamide 0.012 per cent (42.65 per cent) showed poor efficacy with less than 47 per cent reduction in population. However, all the treatments were significantly superior over control in bringing down the population of *S. dorsalis* on chillies at one day after spraying.

The mean data recorded at three days after treatment showed that the most effective treatment was spinosad 0.015 per cent with 93.15 per cent reduction in population of *S. dorsalis* over untreated check and was significantly superior over all the other treatments. The next best treatment was diafenthiuron 0.045 per cent with 91.66 per cent reduction in population and was significantly superior to the rest of the treatments. The treatments pymetrozine 0.02 per cent (86.98 per cent) and fipronil 0.01 per cent (86.33 per cent) being on par with more than 86 per cent reduction in population and significantly superior over untreated check. The other treatments that followed in the descending order of efficacy were imidacloprid 0.005 per cent, chlorfenapyr 0.01 per cent, clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003 per cent with 75.17, 67.08, 62.21, 58.98 and 57.26 per cent reduction in thrips population over untreated check, respectively. Among the treatments, indoxacarb 0.015 per cent (52.03 per cent) and flubendiamide 0.012 per cent (46.54 per cent) were found to be the least effective with less than 53 per cent mean reduction in population over untreated check. However, all the treatments were effective and significant in reducing the population of *S. dorsalis* over control at three days after spraying.

The mean data at seven days after sprayings showed that among the treatments, spinosad 0.015 per cent was the most effective followed by diafenthiuron 0.045 per cent, pymetrozine 0.02 per cent and fipronil 0.01 per cent with 87.03, 84.88, 78.96 and 77.17 per cent reduction over untreated check. The other treatments that followed in the descending order of efficacy were chlorfenapyr 0.01 per cent, imidacloprid 0.005 per cent, clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003 per cent with 69.20, 66.03, 60.20, 57.93 and 56.06 per cent reduction in thrips population over untreated check, respectively. Among the treatments, indoxacarb 0.015 per cent (50.11 per cent) and flubendiamide 0.012 per cent (45.09 per cent) showed poor efficacy with less than 51 per cent reduction in population over control. However, all the treatments were effective and significantly superior over the untreated control in bringing down the population of *S. dorsalis* at seven days after spraying.

The mean data at fourteen days after spraying showed that among the treatments, diafenthiuron 0.045 per cent was the most effective followed by spinosad 0.015 per cent, pymetrozine 0.02 per cent and fipronil 0.01 per cent with 67.86, 67.38, 63.91 and 62.09 per cent reduction over untreated check. The next best treatments were imidacloprid 0.005 per cent (47.22 per cent) and chlorfenapyr 0.01 per cent (46.22 per cent) being on par with more than 46 per cent reduction in population over untreated check. The other treatments that followed in the descending order of efficacy were clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003 per cent with 41.82, 39.93 and 37.14 per cent reduction in thrips population over untreated check, respectively. Among the treatments, indoxacarb 0.015 per cent (35.22 per cent) and flubendiamide 0.012 per cent (32.59 per cent) were found to be the least effective with less than 36 per cent mean reduction in population over untreated check. However, all the treatments were effective and significantly superior over the untreated control in bringing down the population of *S. dorsalis* at fourteen days after spraying.

The data pertaining to the overall cumulative mean efficacy of the treatments during both the seasons revealed that among the treatments, spinosad 0.015 per cent was the most effective followed by diafenthiuron 0.045 per cent, pymetrozine 0.02 per cent and fipronil 0.01 per cent with 82.35, 81.26, 76.97 and 75.62 per cent reduction over untreated check. The other treatments that followed in the descending order of efficacy were imidacloprid 0.005 per cent, chlorfenapyr 0.01 per cent, clothianidin 0.003 per cent, vertimec 0.001 per cent and emamectin benzoate 0.003 per cent with 63.83, 60.32, 55.31, 53.13 and 50.62 per cent reduction in thrips population over untreated check, respectively. Among the treatments,

indoxacarb 0.015 per cent (46.07 per cent) and flubendiamide 0.012 per cent (41.72 per cent) were found to be the least effective with less than 47 per cent reduction in population over control. However, all the treatments were significantly superior over the untreated control in reducing the population of *S. dorsalis* on chillies.

### **EFFECT OF TREATMENTS ON YIELD OF GREEN CHILLIES**

#### ***Kharif2008-2009***

The data pertaining to the yield of chilli presented in Table 4 indicated that the yield obtained from all the plots ranged between 5.04 to 16.10 kg/plot including untreated check. Among the treatments spinosad recorded the highest yield of 16.10 kg/plot (8942.26 kg/ha) with an increase of 219.37 per cent yield over untreated check and was found significantly superior over all the other treatments. Treatments that were moderately better in recording the higher yields were diafenthiuron (15.88 kg/plot or 8822.28 kg/ha), pymetrozine (14.83 kg/plot or 8236.36 kg/ha), fipronil (14.44 kg/plot or 8024.88 kg/ha), imidacloprid (13.91 kg/plot or 7728.24 kg/ha), chlorfenapyr (13.52 kg/plot or 7512.13 kg/ha), clothianidin (13.26 kg/plot or 7365.22 kg/ha), vertimec (12.80 kg/plot or 7113.12 kg/ha) and emamectin benzoate (10.42 kg/plot or 5788.24 kg/ha) being on par with each other resulting in 215.08, 194.16, 186.60, 176.01, 168.29, 163.04, 154.04 and 106.72 per cent increase in yield respectively over untreated check. Among the treatments, indoxacarb (9.56 kg/plot or 5313.14 kg/ha) and flubendiamide (8.97 kg/plot or 4982.66 kg/ha) were on par with each other and recorded significantly less yield with 89.76 and 77.95 per cent increase in yield but was significantly superior over untreated check. The untreated control recorded the lowest yield (5.04 kg/plot or 2800 kg/ha) and was found inferior to all other treatments. However, from the point of cost benefit ratio (CBR) highest CBR 1: 2.20 was recorded in spinosad followed by diafenthiuron (1: 2.19), fipronil (1: 2.00), imidacloprid (1: 1.93), chlorfenapyr (1: 1.87), clothianidin (1: 1.85), vertimec (1: 1.76), emamectin benzoate (1: 1.42), indoxacarb (1: 1.30) and flubendiamide (1: 1.22).

#### ***Kharif2009-2010***

The data pertaining to the yield of chilli presented in Table 5 indicated that the yield obtained from all the plots ranged between 5.44 to 16.49 kg/plot including untreated control. Among the treatments spinosad recorded the highest yield of 16.49 kg/plot (9162.31 kg/ha) with an increase of 203.38 per cent yield over control and was found significantly superior over all the other treatments. Treatments that were moderately better in recording the higher yields were diafenthiuron (16.28 kg/plot or 9042.33 kg/ha), pymetrozine (15.22 kg/plot or 8456.41 kg/ha), fipronil (14.84 kg/plot or 8244.93 kg/ha), imidacloprid (14.31 kg/plot or 7948.29 kg/ha), chlorfenapyr (13.92 kg/plot or 7732.18 kg/ha), clothianidin (13.65 kg/plot or 7585.27 kg/ha), vertimec (13.20 kg/plot or 7333.17 kg/ha) and emamectin benzoate (10.81 kg/plot or 6008.29 kg/ha) being on par with each other resulting in 199.41, 180.01, 173.01, 163.18, 156.03, 151.16, 142.82 and 98.95 per cent increase in yield respectively over control. Among the treatments, indoxacarb (9.96 kg/plot or 5533.19 kg/ha) and flubendiamide (9.36 kg/plot or 5202.71 kg/ha) were on par with each other and recorded significantly less yield with 83.22 and 72.27 per cent increase in yield but was significantly superior over control. The untreated control recorded the lowest yield (5.44 kg/plot or 3020.05 kg/ha) and was found inferior to all other treatments. However, from the point of cost benefit ratio (CBR) highest CBR 1: 2.23 was recorded in spinosad followed by diafenthiuron (1: 2.21), fipronil (1: 2.03), imidacloprid (1: 1.96), chlorfenapyr (1: 1.90), clothianidin (1: 1.88), vertimec (1: 1.79), emamectin benzoate (1: 1.46), indoxacarb (1: 1.34) and flubendiamide (1: 1.26).

#### ***Mean yield for kharif2008-09 and 09-10***

The mean data pertaining to the yield of chilli presented in Table 6 indicated that the yield obtained from all the plots ranged between 5.24 to 16.29 kg/plot including untreated control. Among the treatments spinosad recorded the highest yield of 16.29 kg/plot (9052.36 kg/ha) with an increase of 211.07 per cent yield over control and was found significantly superior over all the other treatments. Treatments that were moderately better in recording the higher yields were diafenthiuron (16.08 kg/plot or 8932.38 kg/ha), pymetrozine (15.02 kg/plot or 8346.45 kg/ha), fipronil (14.64 kg/plot or 8134.97 kg/ha), imidacloprid (14.11 kg/plot or 7838.33 kg/ha), chlorfenapyr (13.72 kg/plot or 7622.22 kg/ha), clothianidin (13.46 kg/plot or 7475.30 kg/ha), vertimec (13.00 kg/plot or 7223.20 kg/ha) and emamectin benzoate (10.62 kg/plot or 5898.31 kg/ha) being on par with each other resulting in 206.95,

186.81, 179.55, 169.35, 161.93, 156.88, 148.22 and 102.69 per cent increase in yield respectively over control. Among the treatments, indoxacarb (9.76 kg/plot or 5423.21 kg/ha) and flubendiamide (9.17 kg/plot or 5092.73 kg/ha) were on par with each other and recorded significantly less yield with 86.36 and 75.00 per cent increase in yield but was significantly superior over control. The untreated control recorded the lowest yield (5.24 kg/plot or 2910.05 kg/ha) and was found inferior to all other treatments. However, from the point of cost benefit ratio (CBR) highest CBR 1: 2.22 was recorded in spinosad followed by diafenthiuron (1: 2.20), fipronil (1: 2.01), imidacloprid (1: 1.95), chlorfenapyr (1: 1.89), clothianidin (1: 1.86), vertimec (1: 1.78), emamectin benzoate (1: 1.44), indoxacarb (1: 1.32) and flubendiamide (1: 1.24).

The overall cumulative efficacy of the observations made at one, three, ten and fourteen days after each of five sprayings at fourteen days intervals during both the years showed that spinosad was significantly superior over rest of the treatments and showed 82.35 per cent mean reduction of thrips population. This is probably due to its recent introduction in to chilli ecosystem which had novel mode of action. It acts on the central nervous system by activation of nicotinic acetylcholine receptors and also effects on GABA receptor. The second best treatment was found to be diafenthiuron with 81.26 per cent mean reduction of thrips population. This is probably due to diafenthiuron is a new molecule belonging to thiourea group and it inhibits the mitochondrial ATP synthesis. The next best treatment was pymetrozine with 76.97 per cent mean reduction of thrips population. This is probably due to pymetrozine is a new molecule belonging to pyridine azomethines group and it blocks the stylet penetration. The next best treatment was found to be fipronil with 75.62 per cent mean reduction of thrips population. This is probably due to fipronil is a new molecule belonging to phenyl pyrazole group. It acts on central nervous system by blocking GABA receptors (Walunj *et al.*, 2000).

### References

- Agricultural statistics at a glance 2009.** <http://www.agricoop.nic.in>
- Ananthkrishnan T N 1971.** Thrips (Thysanoptera) in agriculture, horticulture, forestry - Diagnosis, bionomics and control. *Journal of Scientific and Industrial Research* 30: 113-146.
- Borah D C 1987.** Biology of *Polyphagotarsonemus latus* (Banks) (Acari: Tarsonemidae) and *Scirtothrips dorsalis* (Thysanoptera : Thripidae) infesting chilli and their natural enemies. Ph.D. Thesis submitted to University of Agricultural Sciences, Dharwad, Karnataka.
- FAO (Food and Agricultural Organization) 1979.** FAO methods recommended method for detection and measurement of resistance in agricultural pests to pesticides. *FAO Plant Protection Bulletin* 127:2-5.
- Flemming R and Retnakaran A 1985.** Evaluation of single treatment data using Abbott's formula with reference to insects. *Indian Journal of Economic Zoology* 78: 1179-1181.
- Joia B S, Jaswinder Kaur and Udean A S 2001.** Persistence of ethion residues on/in green chilli. *Proceedings of 2nd National Symposium on Integrated Pest Management (IPM) in Horticultural Crops, New molecules, Biopesticides and Environment.* Bangalore, 17-19 October: 174-175.
- Kranthi K R, Jadhav D R, Kranthi S, Wanjari R R, Ali S S and Russell D 2002.** Insecticide resistance in five major insect pests of cotton in India. *Crop Protection* 21: 449-460.
- Prasad D K, Madhumathi T, Arjunarao P and Srinivasarao V 2008.** Status of OP and Pyrethroid resistance in *Spodoptera litura* (Fab). on cotton. *Pesticide Research Journal* 20 (1):103-106.
- Rao D and Ahmed K 1985.** Evaluation of certain insecticides for the control of the pest complex on chilli in Andhra Pradesh. *Pesticides* 19(2): 41-44.
- Reddy G P V, Prasad V D and Rao R S 1992.** Relative resistance in chilli thrips, *Scirtothrips dorsalis* (Hood) populations in Andhra Pradesh to some conventional insecticides. *Indian Journal of Plant Protection* 20(2): 218-222.



**Varadharajan S 1994.** Studies on host plant resistant and biology of chilli thrips, *Scirtothrips dorsalis* Hood (Thysanoptera: Thripidae). M.Sc. (Ag.) Thesis submitted to Annamalai University, Annamalainagar pp.150.

**Wang K Y, Jiang X Y, Yi M Q, Chen B K and Xia X M 2002.** Insecticide resistance of *Spodoptera exigua*. Acta – Phytophylacica – Sinica 29(3): 229-234.

Table 1: Mean Efficacy of treatments against *S. dorsalis* on chilli during *kharif* 2008-2009

Treatments	Conc. (%)	PTC (Mean no. of thrips/ leaf)	Mean per cent reduction over control				
			1 DAS	3 DAS	7 DAS	14 DAS	Overall
Fipronil	0.01	4.29	77.48 (61.68) <sup>d</sup>	86.67 (68.61) <sup>c</sup>	77.52 (61.71) <sup>d</sup>	62.40 (52.18) <sup>c</sup>	76.02 (60.69) <sup>d</sup>
Pymetrozine	0.02	4.00	78.64 (62.49) <sup>c</sup>	87.32 (69.18) <sup>c</sup>	79.32 (62.96) <sup>c</sup>	64.22 (53.26) <sup>b</sup>	77.38 (61.61) <sup>c</sup>
Imidacloprid	0.005	4.43	67.52 (55.26) <sup>e</sup>	75.51 (60.35) <sup>d</sup>	66.38 (54.57) <sup>f</sup>	47.53 (43.58) <sup>d</sup>	64.24 (53.28) <sup>e</sup>
Diafenthiuron	0.045	4.10	81.26 (64.37) <sup>b</sup>	92.00 (73.63) <sup>b</sup>	85.23 (67.43) <sup>b</sup>	68.17 (55.66) <sup>a</sup>	81.66 (64.66) <sup>b</sup>
Vertimec	0.001	4.93	56.29 (48.61) <sup>g</sup>	59.32 (50.37) <sup>g</sup>	58.29 (49.77) <sup>h</sup>	40.24 (39.37) <sup>f</sup>	53.53 (47.03) <sup>h</sup>
Spinosad	0.015	3.58	82.43 (65.24) <sup>a</sup>	93.49 (75.31) <sup>a</sup>	87.38 (69.23) <sup>a</sup>	67.69 (55.36) <sup>a</sup>	82.75 (65.48) <sup>a</sup>
Emamectin benzoate	0.003	5.05	52.61 (46.50) <sup>h</sup>	57.60 (49.38) <sup>h</sup>	56.41 (48.68) <sup>i</sup>	37.45 (37.73) <sup>g</sup>	51.02 (45.58) <sup>i</sup>
Indoxacarb	0.015	5.11	47.52 (43.58) <sup>i</sup>	52.37 (46.36) <sup>i</sup>	50.46 (45.26) <sup>j</sup>	35.53 (36.59) <sup>h</sup>	46.47 (42.97) <sup>j</sup>
Flubendiamide	0.012	5.16	43.25 (41.12) <sup>j</sup>	46.88 (43.21) <sup>j</sup>	45.45 (42.39) <sup>k</sup>	32.90 (35.00) <sup>i</sup>	42.12 (40.46) <sup>k</sup>
Clothianidin	0.003	4.98	57.60 (49.37) <sup>g</sup>	62.55 (52.27) <sup>f</sup>	60.55 (51.09) <sup>g</sup>	42.13 (40.47) <sup>e</sup>	55.71 (48.28) <sup>g</sup>
Chlorfenapyr	0.01	4.85	59.38 (50.41) <sup>f</sup>	67.42 (55.20) <sup>e</sup>	69.56 (56.52) <sup>e</sup>	46.53 (43.01) <sup>d</sup>	60.72 (51.19) <sup>f</sup>
Untreated check	-	8.44	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
F-test		NS	Sig.	Sig.	Sig.	Sig.	Sig.
SEd		-	0.39	0.45	0.35	0.33	0.30
CD (p=0.05)		-	0.80	0.92	0.72	0.69	0.63

Figures in parentheses are angular transformed values

Numbers followed by same letter in each column are not significantly different

Table 2: Mean Efficacy of treatments against *S. dorsalis* on chilli during *kharif* 2009-2010

Treatments	Conc. (%)	PTC (Mean no. of thrips/ leaf)	Mean per cent reduction over control				
			1 DAS	3 DAS	7 DAS	14 DAS	Overall
Fipronil	0.01	4.08	76.27 (60.89) <sup>b</sup>	85.99 (68.09) <sup>c</sup>	76.81 (61.25) <sup>c</sup>	61.78 (51.82) <sup>b</sup>	75.21 (60.17) <sup>b</sup>
Pymetrozine	0.02	4.12	77.44 (61.69) <sup>b</sup>	86.64 (68.64) <sup>c</sup>	78.61 (62.50) <sup>c</sup>	63.60 (52.91) <sup>b</sup>	76.57 (61.09) <sup>b</sup>
Imidacloprid	0.005	4.13	66.32 (54.55) <sup>c</sup>	74.83 (59.92) <sup>d</sup>	65.67 (54.15) <sup>e</sup>	46.91 (43.23) <sup>c</sup>	63.43 (52.81) <sup>c</sup>
Diafenthiuron	0.045	3.88	80.06 (63.53) <sup>a</sup>	91.32 (73.03) <sup>b</sup>	84.52 (66.92) <sup>b</sup>	67.55 (55.30) <sup>a</sup>	80.86 (64.12) <sup>a</sup>
Vertimec	0.001	4.11	55.08 (47.92) <sup>e</sup>	58.64 (49.98) <sup>g</sup>	57.58 (49.36) <sup>g</sup>	39.62 (39.00) <sup>d</sup>	52.73 (46.57) <sup>f</sup>
Spinosad	0.015	3.71	81.22 (64.39) <sup>a</sup>	92.81 (74.66) <sup>a</sup>	86.67 (68.70) <sup>a</sup>	67.07 (55.00) <sup>a</sup>	81.94 (64.92) <sup>a</sup>
Emamectin benzoate	0.003	4.40	51.41 (45.81) <sup>f</sup>	56.92 (48.98) <sup>g</sup>	55.70 (48.28) <sup>g</sup>	36.83 (37.35) <sup>e</sup>	50.22 (45.12) <sup>g</sup>
Indoxacarb	0.015	4.51	46.32 (42.89) <sup>g</sup>	51.69 (45.97) <sup>h</sup>	49.75 (44.86) <sup>h</sup>	34.91 (36.20) <sup>e</sup>	45.67 (42.51) <sup>h</sup>
Flubendiamide	0.012	4.67	42.05 (40.42) <sup>h</sup>	46.20 (42.82) <sup>i</sup>	44.74 (41.97) <sup>i</sup>	32.28 (34.60) <sup>f</sup>	41.32 (39.99) <sup>i</sup>
Clothianidin	0.003	4.14	56.40 (48.68) <sup>de</sup>	61.87 (51.87) <sup>f</sup>	59.84 (50.68) <sup>f</sup>	41.51 (40.11) <sup>d</sup>	54.90 (47.82) <sup>e</sup>
Chlorfenapyr	0.01	4.31	58.18 (49.71) <sup>d</sup>	66.74 (54.80) <sup>e</sup>	68.85 (56.09) <sup>d</sup>	45.91 (42.65) <sup>c</sup>	59.92 (50.73) <sup>d</sup>
Untreated check	-	8.46	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
F-test		NS	Sig.	Sig.	Sig.	Sig.	Sig.
SEd		-	0.60	0.70	0.60	0.56	0.54
CD (p=0.05)		-	1.25	1.45	1.25	1.17	1.11

Figures in parentheses are angular transformed values

Numbers followed by same letter in each column are not significantly different

Table 3: Mean efficacy of treatments against *S. dorsalis* on chilli during *kharif* 2008-09 and 09-10 (Pooled data)

Treatments	Conc. (%)	PTC (Mean no. of thrips/ leaf)	Mean per cent reduction over control				
			1 DAS	3 DAS	7 DAS	14 DAS	Overall
Fipronil	0.01	4.19	76.87 (61.28) <sup>b</sup>	86.33 (68.35) <sup>c</sup>	77.17 (61.48) <sup>d</sup>	62.09 (52.00) <sup>c</sup>	75.62 (60.42) <sup>c</sup>
Pymetrozine	0.02	4.06	78.04 (62.08) <sup>b</sup>	86.98 (68.90) <sup>c</sup>	78.96 (62.72) <sup>c</sup>	63.91 (53.08) <sup>b</sup>	76.97 (61.35) <sup>b</sup>
Imidacloprid	0.005	4.28	66.92 (54.90) <sup>c</sup>	75.17 (60.13) <sup>d</sup>	66.03 (54.36) <sup>f</sup>	47.22 (43.40) <sup>d</sup>	63.83 (53.04) <sup>d</sup>
Diafenthiuron	0.045	3.99	80.66 (63.94) <sup>a</sup>	91.66 (73.32) <sup>b</sup>	84.88 (67.17) <sup>b</sup>	67.86 (55.48) <sup>a</sup>	81.26 (64.39) <sup>a</sup>
Vertimec	0.001	4.52	55.69 (48.27) <sup>e</sup>	58.98 (50.18) <sup>g</sup>	57.93 (49.57) <sup>h</sup>	39.93 (39.18) <sup>f</sup>	53.13 (46.80) <sup>g</sup>
Spinosad	0.015	3.64	81.83 (64.81) <sup>a</sup>	93.15 (74.97) <sup>a</sup>	87.03 (68.95) <sup>a</sup>	67.38 (55.18) <sup>a</sup>	82.35 (65.19) <sup>a</sup>
Emamectin benzoate	0.003	4.72	52.01 (46.15) <sup>f</sup>	57.26 (49.18) <sup>g</sup>	56.06 (48.48) <sup>i</sup>	37.14 (37.54) <sup>g</sup>	50.62 (45.35) <sup>h</sup>
Indoxacarb	0.015	4.81	46.92 (43.23) <sup>g</sup>	52.03 (46.16) <sup>h</sup>	50.11 (45.06) <sup>j</sup>	35.22 (36.39) <sup>h</sup>	46.07 (42.74) <sup>i</sup>
Flubendiamide	0.012	4.92	42.65 (40.77) <sup>h</sup>	46.54 (43.01) <sup>i</sup>	45.09 (42.18) <sup>k</sup>	32.59 (34.80) <sup>i</sup>	41.72 (40.23) <sup>j</sup>
Clothianidin	0.003	4.56	57.00 (49.03) <sup>e</sup>	62.21 (52.07) <sup>f</sup>	60.20 (50.89) <sup>g</sup>	41.82 (40.29) <sup>e</sup>	55.31 (48.05) <sup>f</sup>
Chlorfenapyr	0.01	4.58	58.78 (50.06) <sup>d</sup>	67.08 (55.00) <sup>e</sup>	69.20 (56.31) <sup>e</sup>	46.22 (42.83) <sup>d</sup>	60.32 (50.96) <sup>e</sup>
Untreated check	-	8.45	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
F-test		NS	Sig.	Sig.	Sig.	Sig.	Sig.
SEd		-	0.48	0.56	0.46	0.43	0.41
CD (p=0.05)		-	1.00	1.17	0.96	0.90	0.85

Figures in parentheses are angular transformed values

Numbers followed by same letter in each column are not significantly different

Table 4: Effect of treatments on yield of green chillies during *kharif* 2008-09

Treatments	Conc. (%)	Kg/plot	Kg/ha	% increase over control	Gross returns Rs/ha	Cost Benefit ratio
Fipronil	0.01	14.44 <sup>b</sup>	8024.88	186.60	160497.60	1:2.00
Pymetrozine	0.02	14.83 <sup>b</sup>	8236.36	194.16	164727.20	-
Imidacloprid	0.005	13.91 <sup>bc</sup>	7728.24	176.01	154564.80	1:1.93
Diafenthiuron	0.045	15.88 <sup>a</sup>	8822.28	215.08	176445.60	1:2.19
Vertimec	0.001	12.80 <sup>d</sup>	7113.12	154.04	142262.40	1:1.76
Spinosad	0.015	16.10 <sup>a</sup>	8942.26	219.37	178845.20	1:2.20
Emamectin benzoate	0.003	10.42 <sup>e</sup>	5788.24	106.72	115764.80	1:1.42
Indoxacarb	0.015	9.56 <sup>ef</sup>	5313.14	89.76	106262.80	1:1.30
Flubendiamide	0.012	8.97 <sup>f</sup>	4982.66	77.95	99653.20	1:1.22
Clothianidin	0.003	13.26 <sup>cd</sup>	7365.22	163.04	147304.40	1:1.85
Chlorfenapyr	0.01	13.52 <sup>cd</sup>	7512.13	168.29	150242.60	1:1.87
Untreated check	-	5.04 <sup>g</sup>	2800.00	-	56000.00	1:0.70
F-test		Sig.				
SEd		0.49				
CD (p=0.05)		1.03				

Numbers followed by same letter in each column are not significantly different

Table 5: Effect of treatments on yield of green chillies during *kharif* 2009-10

Treatments	Conc. (%)	Kg/plot	Kg/ha	% increase over control	Gross returns Rs/ha	Cost Benefit ratio
Fipronil	0.01	14.84 <sup>cd</sup>	8244.93	173.01	164898.60	1:2.03
Pymetrozine	0.02	15.22 <sup>bc</sup>	8456.41	180.01	169128.20	-
Imidacloprid	0.005	14.31 <sup>cde</sup>	7948.29	163.18	158965.80	1:1.96
Diafenthiuron	0.045	16.28 <sup>ab</sup>	9042.33	199.41	180846.60	1:2.21
Vertimec	0.001	13.20 <sup>f</sup>	7333.17	142.82	146663.40	1:1.79
Spinosad	0.015	16.49 <sup>a</sup>	9162.31	203.38	183246.20	1:2.23
Emamectin benzoate	0.003	10.81 <sup>g</sup>	6008.29	98.95	120165.80	1:1.46
Indoxacarb	0.015	9.96 <sup>gh</sup>	5533.19	83.22	110663.80	1:1.34
Flubendiamide	0.012	9.36 <sup>h</sup>	5202.71	72.27	104054.20	1:1.26
Clothianidin	0.003	13.65 <sup>ef</sup>	7585.27	151.16	151705.40	1:1.88
Chlorfenapyr	0.01	13.92 <sup>def</sup>	7732.18	156.03	154643.60	1:1.90
Untreated check	-	5.44 <sup>i</sup>	3020.05	-	60401.00	1:0.75
F-test		Sig.				
SEd		0.53				
CD (p=0.05)		1.10				

Numbers followed by same letter in each column are not significantly different

Table 6: Effect of treatments on yield of green chillies during kharif 2008-09 and 2009-10 (pooled data)

Treatments	Conc. (%)	Kg/plot	Kg/ha	% increase over control	Gross returns Rs/ha	Cost Benefit ratio
Fipronil	0.01	14.64 <sup>bc</sup>	8134.97	179.55	162699.40	1:2.01
Pymetrozine	0.02	15.02 <sup>b</sup>	8346.45	186.81	166929.04	-
Imidacloprid	0.005	14.11 <sup>cd</sup>	7838.33	169.35	156766.55	1:1.95
Diafenthiuron	0.045	16.08 <sup>a</sup>	8932.38	206.95	178647.53	1:2.20
Vertimec	0.001	13.00 <sup>e</sup>	7223.20	148.22	144464.06	1:1.78
Spinosad	0.015	16.29 <sup>a</sup>	9052.36	211.07	181047.15	1:2.22
Emamectin benzoate	0.003	10.62 <sup>f</sup>	5898.31	102.69	117966.24	1:1.44
Indoxacarb	0.015	9.76 <sup>fg</sup>	5423.21	86.36	108464.17	1:1.32
Flubendiamide	0.012	9.17 <sup>g</sup>	5092.73	75.00	101854.51	1:1.24
Clothianidin	0.003	13.46 <sup>de</sup>	7475.30	156.88	149506.10	1:1.86
Chlorfenapyr	0.01	13.72 <sup>de</sup>	7622.22	161.93	152444.32	1:1.89
Untreated check	-	5.24 <sup>h</sup>	2910.05	-	58200.97	1:0.73
F-test		Sig.				
SEd		0.43				
CD (p=0.05)		0.88				

Numbers followed by same letter in each column are not significantly different